

Black Extended Objects, Naked Singularities and P-Branes

B. Harms and Y. Leblanc

*Department of Physics and Astronomy, The University of Alabama**Box 870324, Tuscaloosa, AL 35487-0324*

We treat the horizons of charged, dilaton black extended objects as quantum mechanical objects. We show that the S matrix for such an object can be written in terms of a p-brane-like action. The requirements of unitarity of the S matrix and positivity of the p-brane tension equivalent severely restrict the number of space-time dimensions and the allowed values of the dilaton parameter a . Generally, black objects transform at the extremal limit into p-branes.

PACS numbers: 4.60.+n, 11.17.+y, 97.60.lf

In Ref. [1] we analyzed the scattering of a massless, point-like particle from a charged, dilatonic black hole [2], assuming the horizon can be regarded as a quantum mechanical object. Following the approach of 't Hooft [3], we showed that under rather general assumptions about quantum gravitodynamics, the S matrix can be written in terms of a “string-like” action with an imaginary string tension depending upon the dilaton parameter a . For two values of a ($a = \sqrt{3/5}, \sqrt{7}$) the string tension becomes positive real beyond the extremal limit (coincidence of the inner and outer horizons), and black holes undergo a transition to strings in the region where, classically, they would be naked singularities. We proposed this as a new way of implementing the “cosmic censorship” hypothesis and speculated that such transitions could explain the observed extragalactic gamma ray bursts, the gamma rays being the massless excitations of the strings produced at the end point of the gravitational collapse.

In this paper we extend the work of Ref. [1] to higher dimensions by considering black extended objects. Horowitz and Strominger [4] have obtained the metrics for charged black (10-D)-brane solutions of the field equations which extremize the 10-dimensional action

$$S = \frac{1}{16\pi} \int d^{10}x \sqrt{-g} [e^{-2\Phi} [R + 4(\partial\Phi)^2] - \frac{2}{(D-2)!} e^{2\mu\Phi} F^2], \quad (1)$$

where Φ is the dilaton field, R is the scalar curvature of the 10-dimensional space-time, and the field $F_{\mu_1 \dots \mu_{D-2}}$ is a (D-2)-form satisfying $dF = 0$. The metrics for these black p-branes were obtained by requiring rotational and translational symmetry in (10-D) dimensions and then substituting a general ten-dimensional metric with these symmetries into Eq.(1) to obtain an effective D-dimensional action. The goal of finding black (10-D)-brane solutions of Eq.(1) is thus achieved by finding black hole solutions to the D-dimensional action. Gibbons and Maeda [5] have obtained the black hole solutions for such an action. The most general line element associated with static, spherically symmetric solutions to the D-dimensional equations of motion which are asymptotically flat and have a regular horizon is of the form

$$ds^2 = -\lambda^2 dt^2 + \lambda^{-2} d\hat{r}^2 + R^2 d\Omega_{D-2}^2, \quad (2)$$

where λ and R depend only on $\hat{r}$ and $d\Omega_{D-2}^2$ is the line element on the unit (D-2)-sphere. $\hat{r}$ is related to the usual radius coordinate by $r^{D-4}dr = R^{D-4}d\hat{r}$, λ^2 is given by

$$\lambda^2 = [1 - (\frac{r_+}{r})^{D-3}] [1 - (\frac{r_-}{r})^{D-3}]^{1-\gamma(D-3)} , \quad (3)$$

and R^2 is given by

$$R^2 = r^2 [1 - (r_-/r)^b]^\gamma , \quad (4)$$

with $b = D - 3$. The exponent γ is given by

$$\gamma = \frac{2 a^2 (D - 2)}{(D - 3)[2(D - 3) + a^2 (D - 2)]} , \quad (5)$$

with

$$a = [4\mu^2 + 2\mu(7 - D) + 2\frac{D - 1}{(D - 2)}]^{1/2} . \quad (6)$$

The parameters r_- and r_+ are respectively the inner and outer radii of the horizons of a charged black hole. They are related to the mass M and charge Q by

$$\begin{aligned} M &= \frac{1}{2} [1 - (D - 3)\gamma] r_-^{D-3} + r_+^{D-3} \\ Q &= [\frac{\gamma (D - 3)^3 (r_+ r_-)^{D-3}}{2a^2}]^{1/2} . \end{aligned} \quad (7)$$

To calculate the phase shift of a massless particle scattering from an extended black object we follow the procedure of Ref. [6]. The metric of space-time surrounding such an object with a massless particle falling in from spherical angle Ω' can be represented by gluing together two solutions of Einstein's field equations. The gluing is done after a nonconstant shift of one of the Kruskal coordinates,

$$\delta v(\Omega) = p_{in} f(\Omega, \Omega') \quad (8)$$

along the null surface defined by setting the other Kruskal coordinate to zero ($u = 0$). The incident particle's momentum, p_{in} , is measured with respect to the Kruskal coordinates and the Green function f satisfies the equation, at $u = 0$,

$$\frac{A}{g}\Delta f - \frac{g_{,uv}}{g}f = 32\pi A^2 \delta^2(\Omega, \Omega')|_{u=0} , \quad (9)$$

where Δ is the angular Laplacian in D-2 dimensions. The Kruskal coordinates are chosen such that the metric is of the form

$$d\hat{s} = 2 A(u, v)dudv + g(u, v) d\Omega_{D-2}^2 . \quad (10)$$

The conditions for the matching of the two solutions to the field equations after the shifting are

$$A_{,v}|_{u=0} = g_{,v}|_{u=0} = 0 . \quad (11)$$

The forms for $A(u, v)$ and $g(u, v)$ are determined by obtaining the transformations from the $(\hat{r}, t)$ coordinates to the Kruskal coordinates (u, v) . This is accomplished by defining the “tortoise” coordinate ξ as,

$$\xi = \int \frac{d\hat{r}}{\lambda^2} \quad (12)$$

with λ^2 being given by Eq.(3). The Kruskal coordinates are obtained as in, for example, Ref. [7]. The form of the Kruskal coordinates is the same as in the D = 4 case,

$$u = e^{\alpha\xi} e^{\alpha t} \quad (13)$$

$$v = -e^{\alpha\xi} e^{-\alpha t} ,$$

where α is a constant whose value is determined by the matching conditions (Eq.(11)).

These expressions for u and v give for $A(u, v)$ and $g(u, v)$,

$$A(u, v) = -\frac{\lambda^2 e^{-2\alpha\xi}}{2 \alpha^2} \quad (14)$$

$$g(u, v) = R^2 .$$

The vanishing of $A_{,v}$ and $g_{,v}$ at $u = 0$ ($r = r_+$) requires α to be,

$$\alpha = \frac{b}{2 r_+} (1 - (r_-/r_+)^b)^{1-\gamma(D-2)/2} . \quad (15)$$

The Green function f which determines the phase shift of the particle being scattered is determined by multiplying through Eq.(9) by g and setting,

$$\Gamma = \frac{g_{,uv}}{A} \Big|_{u=0} = \frac{b}{2} (1 - (r_-/r_+)^b) [2 + \gamma b (r_-/r_+)^b (1 - (r_-/r_+)^b)^{-1}] . \quad (16)$$

To simplify the equation determining f we take the “north pole” as the direction of incidence, giving for f ,

$$\Delta f - \Gamma f = -2 \pi \kappa \delta(\theta), \quad (17)$$

with

$$\kappa = -16 Ag|_{u=0} = \frac{32 e^{-1} r_+^4}{b^2} [1 - (r_-/r_+)^b]^{2\gamma-1} , \quad (18)$$

where the constant of integration has been chosen such that this expression for κ agrees with that obtained for the Schwarzschild black hole ($D = 4$ and $\gamma = 0$).

The solution to Eq. (17) has the same form as in [1]. After expanding the δ -function in spherical harmonics and determining the expansion coefficients, we find

$$f = \kappa \sum_l \frac{l + \frac{1}{2}}{l(l+1) + \Gamma} P_l(\cos \theta) , \quad (19)$$

where Γ is as defined in Eq.(16) and $P_l(\cos \theta)$ is the l -th order Legendre polynomial.

Having the function f we can calculate the scattering matrix [3,6],

$$< u(\Omega) | v(\Omega) > = N \exp \left(i \int f^{-1}(\Omega, \Omega') v(\Omega') u(\Omega) \right) , \quad (20)$$

where N is a normalization constant and the propagator is given by

$$f^{-1} = \frac{(\Gamma - \Delta)}{2\pi \kappa} . \quad (21)$$

Fourier transforming the S matrix to the momentum representation we find

$$\begin{aligned}
\langle p_{out}(\Omega) | p_{in}(\Omega) \rangle &= N' \int Du(\Omega) \int Dv(\Omega) \\
&\times \exp \left[\int d^{D-2} \Omega \left(\frac{i}{2\pi \kappa} (\Gamma u v + \partial_\Omega v \partial_\Omega u) + i u p_{out} - i v p_{in} \right) \right].
\end{aligned} \tag{22}$$

The “mass” term (Γ term), which is due to the curvature of the horizon, will be neglected on the assumption that a patch can be chosen small enough that it is locally flat. To write Eq.(22) as a covariant expression we introduce the coordinates

$$x^0 = (v + u)/2 ; x_{D-1} = (v - u)/2 , \tag{23}$$

where the metric $x^2 = \mathbf{x}^2 - x_0^2$ is used. In terms of these coordinates we can write Eq.(22) as,

$$\begin{aligned}
\langle p_{out}(\Omega) | p_{in}(\Omega) \rangle &= C \int Dx^\mu(\sigma) Dg^{ab} \\
&\times \exp \left[\int d^{D-2} \sigma \left(\frac{-T}{2} [\sqrt{g} g^{ab} \partial_a x^\mu \partial_b x^\mu + i x^\mu p^\mu(\sigma)] \right) \right],
\end{aligned} \tag{24}$$

where the integration $d^{D-2} \sigma$ is over variables which are orthogonal to the membrane coordinates generated by x_0 and x_{D-1} . The string tension equivalent is given by

$$T = \frac{i}{\pi \kappa} . \tag{25}$$

The action term in Eq.(24) is of the Polyakov form except for the factor of i , which prevents this S matrix element from satisfying unitarity. To overcome this difficulty we can analytically continue to the region in parameter space where $r_- > r_+$ and require that the exponent $2\gamma - 1$ in Eq.(18) satisfy the condition

$$2\gamma - 1 = \frac{n}{2} , \quad n = odd , \tag{26}$$

where γ is given in Eq.(5). The values of a^2 are thus

$$a^2 = \frac{2(D-3)^2 (1 + n/2)}{(D-2)(7-D) - (D-3)(D-2) n/2} . \tag{27}$$

If we require a to be real and the p-brane tension to be real and positive, then only $D = 4, 5$ are allowed. The values of a for $D = 4$ were determined in Ref. [1] to be $a = \sqrt{3/5}, \sqrt{7}$ for

$n = 1, 5$ respectively. For the $D = 5$ case only $n = 1$ is allowed, and the corresponding value of the dilaton parameter is $a = 2$. For these values of a the constant, T , is real and positive, and the action is that of a string theory for $D = 4$ and that of a membrane theory in $D = 5$.

If we allow T to be negative, then acceptable values of a are obtained for $D = 4, 5, \dots, 10$. For $D > 5$ however only $n = -1$ leads to acceptable values of a . For a string T is the energy per unit length of the string. So a negative value of T does not seem to have any physical meaning.

If we had carried through the curvature term in Eq.(22), we could have used the condition $\Gamma = 0$, which gives the relation

$$\left(\frac{r_-}{r_+}\right)^{D-3} = \frac{a^2 (D-2) + 2(D-3)}{2(D-3)} . \quad (28)$$

From Eq.(7) we also have

$$\left(\frac{r_-}{r_+}\right)^b = \frac{M^2 \left[1 + \sqrt{1 - \frac{8a^2 Q^2 \delta^2}{\gamma b^3 M^2} [1 - \gamma b]}\right]^2 \gamma b^3}{\delta^2 8a^2 Q^2 [1 - \gamma b]^2} , \quad (29)$$

where we have used the definitions

$$\begin{aligned} b &= D - 3 , \\ \delta &= \frac{(D-2) \pi^{(D-3)/2}}{8\Gamma(\frac{D-1}{2})} . \end{aligned} \quad (30)$$

Using Eqs.(28) and (29) we can obtain the ratio Q/M in terms of a and D for which the curvature term vanishes.

As discussed in [1] the picture we obtain from the foregoing analysis is that the charged, dilaton extended black objects avoid the classical naked singularity for the case $r_- > r_+$ by undergoing a phase transition at the extremal point ($r_- = r_+$) and becoming p-branes for the values of the dilaton parameter given above.

Our analysis shows that the results of Refs. [1,3] can be extended to the case of charged, dilaton extended black objects. Using the constraints of unitarity and the positivity of the string tension, we find that the only acceptable values of the dilaton parameter a are $a = \sqrt{3/5}, \sqrt{7}$ for $D = 4$ and $a = 2$ for $D = 5$. Since $D = 4, 5$ are the only allowed

values of D , the only acceptable solutions to Eq.(1) are the black 6-brane and the black 5-brane, which undergo transitions into strings and membranes respectively at the extremal limit. The conditions of unitarity of the S matrix and positivity of the p-brane tension put surprisingly strong restrictions on the values of D and a . Furthermore duality in the dilaton parameter as defined in Ref. [1] exists only in 4 dimensions.

ACKNOWLEDGMENTS

This research was supported in part by the U.S. Department of Energy under Grant No. DE-FG05-84ER40141.

REFERENCES

- [1] P.H. Cox, B. Harms and Y. Leblanc, University of Alabama preprint, UAHEP-9213.
- [2] D. Garfinkle, G. Horowitz and A. Strominger, Phys. Rev. **D43**, 3140(1991).
- [3] G. 't Hooft, Nuc. Phys. **335**, 138(1990); See also H. Verlinde and E. Verlinde, Princeton preprint PUPT-1279(1991).
- [4] G. Horowitz and A. Strominger, Nuc. Phys. **B360**, 197(1991).
- [5] G. Gibbons and K. Maeda, Nuc. Phys. **B298**, 741(1988).
- [6] T. Dray and G. 't Hooft, Nuc. Phys. **B253**, 173(1985).
- [7] R. Adler, M. Bazin and M. Shiffer, *Introduction to General Relativity*, McGraw-Hill, New York (1965).